

# Solomon Practice Paper

## Further Pure Mathematics 3F

Time allowed: 90 minutes

Centre: [www.CasperYC.club](http://www.CasperYC.club)

Name:

Teacher:

Question	Points	Score
1	6	
2	8	
3	11	
4	11	
5	11	
6	13	
7	15	
Total:	75	

How I can achieve better:

- 
- 
- 



Last updated:

December 24, 2025







3. A transformation  $T$  from the  $z$ -plane to the  $w$ -plane is defined by

$$w = \frac{z + 2\mathbf{i}}{z - \mathbf{i}}, \quad z \neq \mathbf{i},$$


where  $z = x + \mathbf{i}y$ ,  $w = u + \mathbf{i}v$  and  $x, y, u$  and  $v$  are real.

- (a) Show that the circle  $|z| = 1$  is mapped onto a straight line in the  $w$ -plane under  $T$  and find an equation of the line. [5]

The circle  $|z - (a + \mathbf{i}b)| = r$  in the  $z$ -plane is mapped under  $T$  onto the circle  $|w| = 2$  in the  $w$ -plane, where  $a, b$  and  $r$  are real.

- (b) Find the values of  $a, b$  and  $r$ . [6]

Total: 11



4. The points  $A, B$  and  $C$  with coordinates  $(x_{-1}, y_{-1}), (x_0, y_0)$  and  $(x_1, y_1)$  respectively lie on the curve  $y = f(x)$  with  $x_1 - x_0 = x_0 - x_{-1} = h$ .

(a) Use the first three terms of the Taylor series expansion in ascending powers of  $(x - x_0)$  to show that [5]

$$\left(\frac{d^2y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{h^2}.$$

The variable  $y$  satisfies the differential equation

$$\frac{d^2y}{dx^2} + (x + 2)\frac{dy}{dx} - 3y = 0 \quad \text{with } y = 1 \text{ at } x = 0 \text{ and } y = 1.2 \text{ at } x = 0.1.$$

(b) Use the approximations


$$\left(\frac{dy}{dx}\right)_0 \approx \frac{y_1 - y_{-1}}{2h}$$

and

$$\left(\frac{d^2y}{dx^2}\right)_0 \approx \frac{y_1 - 2y_0 + y_{-1}}{h^2}$$

with a step length of 0.1 to estimate the value of  $y$  at  $x = 0.2$ .

Total: 11







7. The plane  $\Pi_1$  has vector equation

$$\mathbf{r} = 3\mathbf{i} + \mathbf{j} - 4\mathbf{k} + \lambda(\mathbf{j} + 2\mathbf{k}) + \mu(\mathbf{i} + \mathbf{j} + \mathbf{k}).$$

- (a) Find a vector  $\mathbf{n}$  which is normal to  $\Pi_1$ . [3]
- (b) Hence find a vector equation of  $\Pi_1$  in the form  $\mathbf{r} \cdot \mathbf{n} = p$ . [2]
- (c) Find the perpendicular distance between  $\Pi_1$  and the point  $A$  with position vector  $2\mathbf{i} + \mathbf{j} + 4\mathbf{k}$ , giving your answer in the form  $a\sqrt{6}$ , where  $a \in \mathbb{Q}$ . [4]

The plane  $\Pi_2$  has equation  $\mathbf{r} \cdot (\mathbf{i} + b\mathbf{j}) = -4$ .

The angle between  $\Pi_1$  and  $\Pi_2$  is  $30^\circ$ .

- (d) Find the possible values of the constant  $b$ . [6]

Total: 15

